

Vertex Poisson algebras associated with Courant algebroids and their deformations; I

Gaywalee Yamskulna¹²

Department of Mathematical Sciences, Illinois State University, Normal, IL 61790
and

Institute of Science, Walailak University, Nakon Si Thammarat, Thailand

Abstract

This is the first of two papers on vertex Poisson algebras associated with Courant algebroids, and their deformations. In this work, we study relationships between vertex Poisson algebras and Courant algebroids. For any $\mathbb{N}$ -graded vertex Poisson algebra $A = \coprod_{n \in \mathbb{N}} A_{(n)}$, we show that $A_{(1)}$ is a Courant $A_{(0)}$ -algebroid. On the other hand, for any Courant $\mathcal{A}$ -algebroid $\mathcal{B}$, we construct an $\mathbb{N}$ -graded vertex Poisson algebra $A = \coprod_{n \in \mathbb{N}} A_{(n)}$ such that $A_{(0)}$ is $\mathcal{A}$ and the Courant $\mathcal{A}$ -algebroid $A_{(1)}$ is isomorphic to $\mathcal{B}$ as a Courant $\mathcal{A}$ -algebroid.

1 Introduction

A vertex Poisson algebra is an analogue of the notion of a Poisson algebra in the category of vertex algebras. It is a combination of a commutative vertex algebra structure (or equivalently a differential algebra structure) and a vertex Lie algebra structure with a natural compatibility condition (see [BeD, FB]). In fact, one may consider vertex Poisson algebras as the classical limit of vertex algebras (see [BeD, FB]). Important features of the formalism of vertex Poisson algebras can be traced back to the work of I.M. Gelfand, L.A. Dickey and others on the Hamiltonian structure of integrable hierarchies of soliton equations (see [Di]). They play a prominent role in the study of certain connections between the classical and quantum Drinfeld-Sokolov reductions (see [FB] Chapter 15). They also play a crucial role in the study of generating subspaces of vertex algebras with a certain property analogous to the well known Poincaré-Birkhoff-Witt spanning property (see [Li2, Li3] and references therein).

Courant algebroids first appeared in the work of T. Courant on Dirac structures on manifolds (see [Co]). They were later studied by Liu, Weinstein, and Xu who used Courant algebroids to generalize the notion of the Drinfeld double to Lie bialgebroids (see [LiuWX]). One can think of a Courant algebroid as a certain kind of algebroid for which the condition of antisymmetry, the Leibniz rule and the Jacobi identity for the bracket are all relaxed in certain ways. Examples of Courant algebroids are the doubles of Lie bialgebras and the bundle $TM \oplus T^*M$. (see [LiuX, Co]). Courant algebroids play an important role in the study of dynamical r -matrices (see [LiuX]). Also, it was shown in [RoW] that Courant algebroids can be viewed as strongly homotopy Lie algebras. Furthermore, Courant algebroids have a

¹E-mail address: gyamsku@ilstu.edu

²Partially supported by a New Faculty Initiative grant from the College of Arts and Sciences, ISU

close connection to gerbs (see [BrC]) and play a significant role in the study of the theory of vertex algebroids (see [Br]).

The purpose of this paper is to study relationships between Courant algebroids and $\mathbb{N}$ -graded vertex Poisson algebras. Precisely, we show that one can obtain Courant algebroids from $\mathbb{N}$ -graded vertex Poisson algebras and vice versa. For an $\mathbb{N}$ -graded vertex Poisson algebra $A = \coprod_{n \in \mathbb{N}} A_{(n)}$, the homogeneous subspace $A_{(0)}$ is a unital commutative associative algebra and the homogeneous subspace $A_{(1)}$ is a module of an associative algebra $A_{(0)}$. Moreover, the skew symmetry and the half commutator formula of a vertex Poisson algebra give rise to several compatibility relations. These structures on $A_{(0)} \oplus A_{(1)}$ are summarized in the notion of what was called a 1-truncated conformal algebra. By using skew symmetry and the fact that a family of bilinear operations

$$i : A \otimes_{\mathbb{C}} A \rightarrow A, \quad a \otimes b \mapsto a_i b$$

are derivations of the commutative product on A , we show that $A_{(1)}$ is, in fact, a Courant $A_{(0)}$ -algebroid. On the other hand, for a given Courant $\mathcal{A}$ -algebroid $\mathcal{B}$, we construct an $\mathbb{N}$ -graded vertex Poisson algebra $A = \coprod_{n \in \mathbb{N}} A_{(n)}$ with $A_{(0)} = \mathcal{A}$ and the homogeneous subspace $A_{(1)}$ isomorphic to $\mathcal{B}$ as a Courant algebroid. For a given 1-truncated conformal algebra $\mathbb{A} \oplus \mathbb{B}$ over $\mathbb{C}$, we set $C(\mathbb{A} \oplus \mathbb{B}) = \mathbb{C}[\mathcal{D}] \otimes (\mathbb{A} \oplus \mathbb{B})$ where $\mathcal{D}$ is a formal variable. We use the fact that $\mathbb{A} \oplus \mathbb{B}$ is a 1-truncated conformal algebra to show that a certain quotient space $\mathcal{C}_{\mathbb{B}}$ of $C(\mathbb{A} \oplus \mathbb{B})$ is a vertex Lie algebra. It was shown in [Li2] that a symmetric algebra of a vertex Lie algebra is a vertex Poisson algebra. Applying this result to $\mathcal{C}_{\mathbb{B}}$, we then obtain a vertex Poisson algebra $S(\mathcal{C}_{\mathbb{B}})$. In fact, $S(\mathcal{C}_{\mathbb{B}})$ is an $\mathbb{N}$ -graded vertex Poisson algebra whose degree-zero subspace is $S(\mathbb{A})$ and whose degree-one subspace is $S(\mathbb{A}) \cdot \mathbb{B}$. Here, $S(\mathbb{A})$ is the symmetric algebra over the space $\mathbb{A}$. For any Courant $\mathcal{A}$ -algebroid $\mathcal{B}$, $\mathcal{A} \oplus \mathcal{B}$ is a 1-truncated conformal algebra. Hence, $S(\mathcal{C}_{\mathcal{B}})$ is an $\mathbb{N}$ -graded vertex Poisson algebra. By using the fact that $\mathcal{B}$ is a Courant $\mathcal{A}$ -algebroid, we can show that a certain quotient space $S_{\mathcal{B}}$ of $S(\mathcal{C}_{\mathcal{B}})$ is an $\mathbb{N}$ -graded vertex Poisson algebra whose the degree-zero subspace is $\mathcal{A}$ and the degree-one subspace is isomorphic to $\mathcal{B}$ as a Courant $\mathcal{A}$ -algebroid.

This paper is organized as follows: In section 2 we recall notions of vertex Lie algebras, and vertex Poisson algebras. In section 3, we review the notions of 1-truncated conformal algebras and Courant $\mathcal{A}$ -algebroids, and we construct Courant algebroids from $\mathbb{N}$ -graded vertex Poisson algebras. In section 4, we construct an $\mathbb{N}$ -graded vertex Poisson algebra from any 1-truncated conformal algebra. In section 5, for any Courant $\mathcal{A}$ -algebroid $\mathcal{B}$, we construct an $\mathbb{N}$ -graded vertex Poisson algebra whose degree-zero subspace is $\mathcal{A}$ and degree-one subspace is isomorphic to $\mathcal{B}$ as a Courant $\mathcal{A}$ -algebroid.

We use here the standard formal variable notations and conventions as defined in [FHL, LLi]. Also, we use a notation $\mathbb{N}$ for the set of nonnegative integers.

2 Vertex Lie algebras, and vertex Poisson algebras

In this section we review the definitions of vertex Lie algebras and vertex Poisson algebras, as well as their properties that will be used in later sections.

Let W be a vector space. Following [P, Li2], for a formal series

$$f(x_1, \dots, x_n) = \sum_{m_1, \dots, m_n \in \mathbb{Z}} u(m_1, \dots, m_n) x_1^{-m_1-1} \dots x_n^{-m_n-1} \in W[[x_1^{\pm 1}, \dots, x_n^{\pm 1}]],$$

we define

$$\text{Sing } f(x_1, \dots, x_n) = \sum_{m_1, \dots, m_n \in \mathbb{N}} u(m_1, \dots, m_n) x_1^{-m_1-1} \dots x_n^{-m_n-1}.$$

For $1 \leq i \leq n$, we have

$$\frac{\partial}{\partial x_i} \text{Sing } f(x_1, \dots, x_n) = \text{Sing } \frac{\partial}{\partial x_i} f(x_1, \dots, x_n). \quad (2.1)$$

For any nonempty subset $S = \{i_1, \dots, i_k\}$ of $\{1, \dots, n\}$, a formal series $f(x_1, \dots, x_n)$ can be viewed as a formal series $f(x_{i_1}, \dots, x_{i_k})$ in variables $x_{i_1}, \dots, x_{i_k}$ with coefficients in the vector space $V[[x_j^{\pm 1} | j \notin S]]$. We define

$$\text{Sing}_{x_{i_1}, \dots, x_{i_k}} f(x_1, \dots, x_n) = \text{Sing } \tilde{f}(x_{i_1}, \dots, x_{i_k}). \quad (2.2)$$

Hence,

$$\text{Sing}_{x_1} f(x_1, \dots, x_n) = \sum_{m \in \mathbb{N}, m_2, \dots, m_n \in \mathbb{Z}} u(m, m_2, \dots, m_n) x_1^{-m-1} x_2^{-m_2-1} \dots x_n^{-m_n-1}$$

and

$$\text{Sing } f(x_1, \dots, x_n) = \text{Sing}_{x_1} \dots \text{Sing}_{x_n} f(x_1, \dots, x_n).$$

Proposition 2.1. [P] Let W be a vector space and let $B \in W((x_1, \dots, x_n))$, $M \in (\text{End } W)[[x_1, \dots, x_n]]$. Then

$$\text{Sing } (M \cdot \text{Sing } (B)) = \text{Sing } (MB).$$

Next, we recall the notions of vertex Lie algebra and vertex Poisson algebra.

Definition 2.2. [K, P] A vertex Lie algebra is a vector space A equipped with a linear operator ∂ and a linear map

$$\begin{aligned} Y_- : A &\rightarrow \text{Hom}(A, x^{-1}A[x^{-1}]) \\ a &\mapsto Y_-(a, x) = \sum_{n \in \mathbb{N}} a_n x^{-n-1} \quad (\text{where } a_n \in \text{End } A) \end{aligned}$$

such that the following axioms hold for $a, b \in A$:

$$a_n b = 0 \quad \text{for } n \text{ sufficiently large}; \quad (2.3)$$

$$Y_-(\partial a, x) = \frac{d}{dx} Y_-(a, x); \quad (2.4)$$

$$Y_-(a, x)b = \text{Sing}(e^{x\partial} Y_-(b, -x)a); \quad (2.5)$$

$$[Y_-(a, x_1), Y_-(b, x_2)] = \text{Sing}(Y_-(Y_-(a, x_1 - x_2)b, x_2)). \quad (2.6)$$

Following [P], we call (2.6) the half commutator formula.

Remark 2.3. In terms of components, (2.4)-(2.6) are equivalent to

$$(\partial a)_n b = -n a_{n-1} b, \quad (2.7)$$

$$a_n b = \sum_{i \geq 0} (-1)^{n+i+1} \frac{1}{i!} \partial^i b_{n+i} a, \quad (2.8)$$

$$a_m b_n c - b_n a_m c = \sum_{i=0}^m \binom{m}{i} (a_i b)_{m+n-i} c, \quad (2.9)$$

for $a, b, c \in A, m, n \in \mathbb{N}$.

Remark 2.4. Any vertex algebra is a vertex Lie algebra with $\partial = \mathcal{D}$. (See appendix for the definition of a vertex algebra.)

Proposition 2.5. [K] For $a \in A$, $[\partial, Y_-(a, x)] = \frac{d}{dx} Y_-(a, x) = Y_-(\partial a, x)$.

Proposition 2.6. [Li2] Let A be a vector space equipped with a linear operator ∂ and equipped with a linear map

$$\begin{aligned} Y_- : A &\rightarrow \text{Hom}(A, x^{-1}A[x^{-1}]) \\ a &\mapsto Y_-(a, x) \end{aligned} \quad (2.10)$$

such that the following conditions hold for all $a, b \in A$:

$$\begin{aligned} [\partial, Y_-(a, x)] &= \frac{d}{dx} Y_-(a, x), \\ Y_-(a, x)b &= \text{Sing}(e^{x\partial} Y_-(b, -x)a). \end{aligned}$$

Then the half commutator formula for an ordered triple (a, b, c) implies the half commutator formula for any permutation of (a, b, c) .

A differential algebra is a commutative associative algebra A with the identity 1 equipped with a derivation ∂ . We often denote the differential algebra by (A, ∂) . A subset U of A generates A as a differential algebra if $\partial^n U$ for $n \in \mathbb{N}$ generate A as an algebra.

Definition 2.7. [FB] A vertex Poisson algebra is a differential algebra (A, ∂) equipped with a vertex Lie algebra structure (Y_-, ∂) (with the same operator ∂) such that for $a, b, c \in A$,

$$Y_-(a, x)(bc) = (Y_-(a, x)b)c + b(Y_-(a, x)c). \quad (2.11)$$

Remark 2.8. In terms of components, (2.11) is equivalent to

$$a_n(bc) = (a_n b)c + b(a_n c) \quad \text{for } a, b, c \in A, n \in \mathbb{N}. \quad (2.12)$$

Here $Y_-(a, x) = \sum_{n \in \mathbb{N}} a_n x^{-n-1}$.

Corollary 2.9. For $a \in A, n \in \mathbb{N}$,

1. a_n is derivation of A , and

2. $Y_-(a, x)1 = 0$, and $Y_-(1, x)a = 0$.

Proof. The second part of 2. follows from the first part of 2., and (2.5). $\square$

A vertex algebra is called *commutative* if $[Y(u, x_1), Y(v, x_2)] = 0$ for all $u, v \in V$. (See appendix for the definition of a vertex algebra.) It was shown in [FHL] that a vertex algebra V is commutative if and only if $u_n v = 0$ for all $u, v \in V$, $n \in \mathbb{N}$.

Proposition 2.10. [Bo, FB, Li1] *If V is a commutative vertex algebra, then V is a commutative associative algebra with the product defined by*

$$u \cdot v = u_{-1}v \quad \text{for } u, v \in V \quad (2.13)$$

and with $\mathbf{1}$ as the identity element. Furthermore, the operator $\mathcal{D}$ of V is a derivation and $Y(u, x)v = (e^{x\mathcal{D}}u)v$ for $u, v \in V$.

Conversely, for any differential algebra (A, ∂) , $(A, Y, 1)$ is a commutative vertex algebra where Y is defined by $Y(a, x)b = (e^{x\partial}a)b$ for $a, b \in A$. This give rise to a canonical isomorphism between the category of commutative vertex algebras and the category of differential algebras. Moreover, a vertex Poisson algebra structure on a vector space A consists of a commutative vertex algebra structure and a vertex Lie algebra structure with a compatibility condition.

Definition 2.11. [Li2] *A $\mathbb{Z}$ -graded vertex Poisson algebra is a vertex Poisson algebra A equipped with a $\mathbb{Z}$ -grading $A = \coprod_{n \in \mathbb{Z}} A_{(n)}$ such that A as an algebra is $\mathbb{Z}$ -graded and such that for $a \in A_{(n)}$, $n, r \in \mathbb{Z}$, $m \in \mathbb{N}$*

$$\begin{aligned} \partial A_{(r)} &\subset A_{(r+1)}, \\ a_m A_{(r)} &\subset A_{(r+n-m-1)}. \end{aligned}$$

An $\mathbb{N}$ -graded vertex Poisson algebra is defined in the obvious way.

Remark 2.12. *Let A be an $\mathbb{Z}$ -graded (respectively, $\mathbb{N}$ -graded) vertex Poisson algebra. Then*

1. $A_{(0)}$ is a commutative associative algebra with the identity 1.
2. For $n \in \mathbb{Z}$ (respectively, $n \in \mathbb{N}$), $A_{(n)}$ is an $A_{(0)}$ -module.
3. $\partial : A_{(0)} \rightarrow A_{(1)}$ is a derivation when we consider $A_{(0)}$ as an algebra and $A_{(1)}$ as an $A_{(0)}$ -module.

Definition 2.13. [Li2] *An ideal of a vertex Poisson algebra A is an ideal I of A as an associative algebra such that*

$$\begin{aligned} \partial I &\subset I \\ a_n I &\subset I \quad \text{for } a \in A, n \in \mathbb{N}. \end{aligned}$$

Corollary 2.14. [Li2]

1. By the half skew symmetry we have $u_n A \subset I$ for $u \in I$, $n \in \mathbb{N}$.

2. Moreover, the quotient space A/I has a natural vertex Poisson algebra structure.

Proposition 2.15. *Let A be a vertex Poisson algebra, and let I be an ideal of A as an associative commutative algebra. Assume that $\partial I \subset I$ and I is generated by W , that is $I = AW$. If $a_n w \in I$ for all $a \in A, w \in W, n \in \mathbb{N}$, then I is an ideal of a vertex Poisson algebra A .*

Proof. Let $a, a' \in A, w \in W, n \in \mathbb{N}$. By (2.12), we have

$$a_n(a'w) = (a_n a')w + a'(a_n w) \in I.$$

It follows that I is an ideal. $\square$

Proposition 2.16. *[Li2, FB] Let R be a vector space equipped with a linear operator $\mathcal{D}$ and let Y_-^0 be a linear map from R to $\text{Hom}(R, x^{-1}R[x^{-1}])$. Denote by $S(R)$ the symmetric algebra over R and we extend $\mathcal{D}$ uniquely to a derivation of $S(R)$. Then Y_-^0 extends to a vertex Poisson algebra structure Y_- on $(S(R), \mathcal{D})$ if and only if $(R, \mathcal{D}, Y_-^0)$ carries the structure of a vertex Lie algebra. Furthermore, such an extension is unique.*

3 From vertex Poisson algebras to Courant algebroids

First, we review the notions of 1-truncated conformal algebras and Courant algebroids. Also, we study the relations between these algebras. Next, we show that for an $\mathbb{N}$ -graded vertex Poisson algebra $V = \coprod_{n \in \mathbb{N}} V_{(n)}$, the homogeneous subspace $V_{(1)}$ is a Courant $V_{(0)}$ -algebroid.

Definition 3.1. *[GMS] A 1-truncated conformal algebra is a graded vector space $C = C_0 \oplus C_1$, equipped with a linear map $\partial : C_0 \rightarrow C_1$ and bilinear operations $(u, v) \mapsto u_i v$ for $i = 0, 1$ of degree $-i - 1$ on $C = C_0 \oplus C_1$ such that the following axioms hold:*

(Derivation) for $a \in C_0, u \in C_1$,

$$(\partial a)_0 = 0; \quad (\partial a)_1 = -a_0; \quad \partial(u_0 a) = u_0 \partial a \quad (3.1)$$

(Commutativity) for $a \in C_0, u, v \in C_1$,

$$u_0 a = -a_0 u; \quad u_0 v = -v_0 u + \partial(v_1 u); \quad u_1 v = v_1 u \quad (3.2)$$

(Associativity) for $\alpha, \beta, \gamma \in C_0 \oplus C_1$,

$$\alpha_0 \beta_i \gamma = \beta_i \alpha_0 \gamma + (\alpha_0 \beta)_i \gamma. \quad (3.3)$$

Remark 3.2.

1. Let $A = \coprod_{n \in \mathbb{N}} A_{(n)}$ be an $\mathbb{N}$ -graded vertex Lie algebra. By equations (2.7)-(2.9), we have that $A_{(0)} \oplus A_{(1)}$ is a 1-truncated conformal algebra.

2. Consequently, for an $\mathbb{N}$ -graded vertex Poisson algebra $A = \coprod_{n \in \mathbb{N}} A_{(n)}$, $A_{(0)} \oplus A_{(1)}$ is a 1-truncated conformal algebra.

A *Leibniz algebra* is a nonassociative algebra Γ satisfying the following condition:

$$u \cdot (v \cdot w) = (u \cdot v) \cdot w + v \cdot (u \cdot w) \quad \text{for } u, v, w \in \Gamma.$$

Any Lie algebra is a Leibniz algebra. In particular, for any vector space W the general linear Lie algebra $\mathfrak{gl}(W)$ is a Leibniz algebra. A *representation of Leibniz algebra* Γ on a vector space W is a Leibniz algebra homomorphism ρ from Γ to $\mathfrak{gl}(W)$.

Let $\mathcal{A}$ be a unital commutative associative algebra (over $\mathbb{C}$). A *Leibniz $\mathcal{A}$ -algebra* is a Leibniz algebra Γ equipped with an $\mathcal{A}$ -module structure. A *module* for a Leibniz $\mathcal{A}$ -algebra Γ is a vector space W equipped with Γ -module structure and an $\mathcal{A}$ -module structure.

Proposition 3.3. [GMS, LiY] *Let $C = C_0 \oplus C_1$ be a graded vector space (over $\mathbb{C}$) equipped with a linear map ∂ from C_0 to C_1 and equipped with bilinear maps $(u, v) \mapsto u_i v$ of degree $-i - 1$ on $C = C_0 \oplus C_1$ for $i = 0, 1$. Then C is a 1-truncated conformal algebra if and only if*

1. The pair $(C_1, [\cdot, \cdot])$ carries the structure of a Leibniz algebra where $[u, v] = u_0 v$ for $u, v \in C_1$.
2. The space C_0 is a C_1 -module with $u \cdot a = u_0 a$ for $u \in C_1, a \in C_0$.
3. The map ∂ is a C_1 -module homomorphism.
4. The subspace ∂C_0 of C_1 annihilates the C_1 -module $C_0 \oplus C_1$.
5. The bilinear map $\langle \cdot, \cdot \rangle$ from $C_1 \otimes C_1$ to C_0 defined by $\langle u, v \rangle = u_1 v$ for $u, v \in C_1$ is a C_1 -module homomorphism and furthermore

$$u_0 a = -a_0 u, \tag{3.4}$$

$$\langle \partial a, u \rangle = -a_0 u, \tag{3.5}$$

$$[u, v] + [v, u] = \partial \langle u, v \rangle, \tag{3.6}$$

$$\langle u, v \rangle = \langle v, u \rangle \tag{3.7}$$

for $a \in C_0, u, v \in C_1$.

Next, we state the definition of a Courant algebroid. We also study the relationship between Courant algebroids and 1-truncated conformal algebras.

Definition 3.4. [Br] *Let $\mathcal{A}$ be a unital commutative associative algebra over $\mathbb{C}$. A Courant $\mathcal{A}$ -algebroid is an $\mathcal{A}$ -module $\mathcal{B}$ equipped with*

1. a structure of Leibniz algebra $[\cdot, \cdot] : \mathcal{B} \otimes_{\mathbb{C}} \mathcal{B} \rightarrow \mathcal{B}$,
2. a homomorphism of Leibniz $\mathcal{A}$ -algebras $\pi : \mathcal{B} \rightarrow \text{Der}(\mathcal{A})$,
3. a symmetric $\mathcal{A}$ -bilinear pairing $\langle \cdot, \cdot \rangle : \mathcal{B} \otimes_{\mathcal{A}} \mathcal{B} \rightarrow \mathcal{A}$,

4. a derivation $\partial : \mathcal{A} \rightarrow \mathcal{B}$ such that $\pi \circ \partial = 0$ which satisfy

$$[u, av] = a[u, v] + \pi(u)(a)v, \quad (3.8)$$

$$\langle [u, v], w \rangle + \langle v, [u, w] \rangle = \pi(u)\langle v, w \rangle, \quad (3.9)$$

$$[u, \partial a] = \partial(\pi(u)a), \quad (3.10)$$

$$\langle u, \partial a \rangle = \pi(u)a, \quad (3.11)$$

$$[u, v] + [v, u] = \partial(\langle u, v \rangle), \quad (3.12)$$

for $a \in \mathcal{A}, u, v, w \in \mathcal{B}$.

Corollary 3.5.

1. $\partial(\mathcal{A})$ annihilates $\mathcal{A}$ and $\mathcal{B}$.
2. $\langle \cdot, \cdot \rangle$ and ∂ are $\mathcal{B}$ -module homomorphisms.

Proof. For 1., since $\pi \circ \partial = 0$, it follows immediately that $\partial(\mathcal{A})$ annihilates $\mathcal{A}$. Next, we will show that $\partial(\mathcal{A})$ annihilates $\mathcal{B}$. Let $a \in \mathcal{A}, u \in \mathcal{B}$. By (3.10)-(3.12), we have

$$[\partial(a), u] = -[u, \partial(a)] + \partial(\langle \partial(a), u \rangle) = -\partial(\pi(u)a) + \partial(\pi(u)a) = 0.$$

2. follows immediately from (3.9), (3.10). $\square$

By the definition of a Courant algebroid, Proposition 3.3, and Corollary 3.5, we have the following.

Proposition 3.6. *Let $\mathcal{A}$ be a unital commutative associative algebra, and let $\mathcal{B}$ be an $\mathcal{A}$ -module. Let $\partial : \mathcal{A} \rightarrow \mathcal{B}$ be a derivation. Then a Courant $\mathcal{A}$ -algebroid structure on $\mathcal{B}$ is exactly equivalent to a 1-truncated conformal algebra structure on $C = \mathcal{A} \oplus \mathcal{B}$ with*

$$\begin{aligned} a_i a' &= 0, \\ u_0 v &= [u, v], \quad u_1 v = \langle u, v \rangle, \\ u_0 a &= \pi(u)(a), \quad a_0 u = -u_0 a \end{aligned}$$

for $a, a' \in \mathcal{A}, u, v \in \mathcal{B}, i = 0, 1$ such that

$$(au)_0 a' = a(u_0 a'), \quad (3.13)$$

$$(au)_1 v = a(u_1 v) = u_1(av), \quad (3.14)$$

$$u_0(av) = a(u_0 v) + (u_0 a)v, \quad (3.15)$$

$$u_0(aa') = a(u_0 a') + (u_0 a)a'. \quad (3.16)$$

Corollary 3.7. *Let e be the identity of $\mathcal{A}$. Then for $u \in \mathcal{B}, u_0 e = 0$.*

Proof. This follows immediately from (3.16). $\square$

Let $(A = \coprod_{n \in \mathbb{Z}} A_{(n)}, \partial)$ be an $\mathbb{N}$ -graded vertex Poisson algebra. By Remark 2.12, we have $A_{(0)}$ is a commutative associative algebra with the identity 1 and $A_{(1)}$ is an $A_{(0)}$ -module. Moreover, $\partial : A_{(0)} \rightarrow A_{(1)}$ is a derivation. By Remark 3.2, we also have that $A_{(0)} \oplus A_{(1)}$ is a 1-truncated conformal algebra.

Theorem 3.8. $A_{(1)}$ is, in fact, a Courant $A_{(0)}$ -algebroid.

Proof. By Proposition 3.6, it is enough to show that (3.13)-(3.16) hold on $A_{(0)} \oplus A_{(1)}$. Let $a, a' \in A_{(0)}, u, v \in A_{(1)}$. By (2.12), we have

$$\begin{aligned} u_0(av) &= a(u_0v) + (u_0a)v, \\ u_0(aa') &= (u_0a)a' + a(u_0a'). \end{aligned}$$

Hence, (3.15), (3.16) hold.

Let $a \in A_{(0)}, u, v \in A_{(1)}$. By (2.12), (3.2), we have

$$\begin{aligned} u_1(av) &= a(u_1v), \text{ and} \\ (au)_1v &= v_1(au) = a(v_1u) = a(u_1v). \end{aligned}$$

Therefore, (3.14) holds.

Next, we will show that (3.13) holds. Let $a, a' \in A_{(0)}, u \in A_{(1)}$. By (2.8), (2.12), (3.2), we have

$$\begin{aligned} (au)_0a' &= \sum_{i \geq 0} (-1)^{i+1} \frac{1}{i!} \partial^i(a'_i(au)) \\ &= -a'_0(au) \\ &= -\{(a'_0a)u + a(a'_0u)\} \\ &= -a(-u_0a') \\ &= a(u_0a'). \end{aligned}$$

Therefore, $A_{(1)}$ is a Courant $A_{(0)}$ -algebroid. $\square$

4 Vertex Poisson algebras associated with 1-truncated conformal algebras

In this section, we construct an $\mathbb{N}$ -graded vertex Poisson algebra from any 1-truncated conformal algebra.

Let $(\mathbb{A} \oplus \mathbb{B}, \partial)$ be a 1-truncated conformal algebra. We set

$$C(\mathbb{A} \oplus \mathbb{B}) = \mathbb{C}[\mathcal{D}] \otimes_{\mathbb{C}} (\mathbb{A} \oplus \mathbb{B}),$$

where $\mathcal{D}$ is a formal variable. We define subspaces $C(\mathbb{A})$ and $C(\mathbb{B})$ in the obvious way. Additionally, we consider $\mathbb{A} \oplus \mathbb{B}$ as a subspace of $C(\mathbb{A} \oplus \mathbb{B})$ via the following map

$$\mathbb{A} \oplus \mathbb{B} \rightarrow C(\mathbb{A} \oplus \mathbb{B}), \quad a + b \mapsto 1 \otimes a + 1 \otimes b.$$

Also, we define a linear operator $\mathcal{D}$ on $C(\mathbb{A} \oplus \mathbb{B})$ by

$$\mathcal{D}(\mathcal{D}^n \otimes u) = \mathcal{D}^{n+1} \otimes u \text{ for } u \in \mathbb{A} \oplus \mathbb{B}, n \in \mathbb{N}, \quad (4.1)$$

and set

$$\hat{\partial} = 1 \otimes \partial - \mathcal{D} \otimes 1 : C(\mathbb{A}) \rightarrow C(\mathbb{A} \oplus \mathbb{B}). \quad (4.2)$$

Next, we define

$$\begin{aligned}\deg(\mathcal{D}^n \otimes a) &= n \quad \text{for } a \in \mathbb{A}, n \in \mathbb{N}, \\ \deg(\mathcal{D}^n \otimes b) &= n + 1 \quad \text{for } b \in \mathbb{B}, n \in \mathbb{N}.\end{aligned}$$

Then $C(\mathbb{A} \oplus \mathbb{B})$ becomes an $\mathbb{N}$ -graded vector space:

$$C(\mathbb{A} \oplus \mathbb{B}) = \coprod_{n \in \mathbb{N}} C(\mathbb{A} \oplus \mathbb{B})_{(n)},$$

where $C(\mathbb{A})_{(0)} = \mathbb{A}$, and for $n \geq 1$

$$\begin{aligned}C(\mathbb{A} \oplus \mathbb{B})_{(n)} &= \mathcal{D}^n \otimes \mathbb{A} \oplus \mathcal{D}^{n-1} \otimes \mathbb{B} \\ &= \{ \mathcal{D}^n \otimes a, \mathcal{D}^{n-1} \otimes b \mid a \in \mathbb{A}, b \in \mathbb{B} \}.\end{aligned}\tag{4.3}$$

The linear map $\hat{\partial}$ is a homogeneous of degree 1 and for $n \geq 1$, we have

$$(\hat{\partial}C(\mathbb{A}))_{(n)} = \{ \hat{\partial}(\mathcal{D}^{n-1} \otimes a) \mid a \in \mathbb{A} \}.$$

Moreover, $\mathcal{D}(\hat{\partial}C(\mathbb{A})) \subset \hat{\partial}C(\mathbb{A})$.

We define a linear map

$$\begin{aligned}Y_-^0 : C(\mathbb{A} \oplus \mathbb{B}) &\rightarrow \text{Hom}(C(\mathbb{A} \oplus \mathbb{B}), x^{-1}C(\mathbb{A} \oplus \mathbb{B})[x^{-1}]) \\ u &\mapsto Y_-^0(u, x) = \sum_{n \in \mathbb{N}} u_n x^{-n-1}\end{aligned}$$

in the following way: for $a, a' \in \mathbb{A}$, $b, b' \in \mathbb{B}$, $u, v \in \mathbb{A} \oplus \mathbb{B}$, $n \geq 1$, $m \in \mathbb{N}$,

$$Y_-^0(a, x)a' = 0, \tag{4.4}$$

$$Y_-^0(a, x)b = a_0 b x^{-1}, \tag{4.5}$$

$$Y_-^0(b, x)a = b_0 a x^{-1}, \tag{4.6}$$

$$Y_-^0(b, x)b' = b_0 b' x^{-1} + b_1 b' x^{-2}, \tag{4.7}$$

$$Y_-^0(u, x)e^{x_1 \mathcal{D}} v = e^{x_1 \mathcal{D}} e^{-x_1 \frac{d}{dx}} Y_-^0(u, x)v, \tag{4.8}$$

$$Y_-^0(\mathcal{D}^n \otimes u, x)\mathcal{D}^m \otimes v = \text{Sing}(e^{x \mathcal{D}} \left(-\frac{d}{dx}\right)^m Y_-^0(v, -x)\mathcal{D}^n \otimes u). \tag{4.9}$$

Remark 4.1. Notice that (4.8) is equivalent to

$$Y_-^0(u, x)\mathcal{D}^n \otimes v = \sum_{i=0}^n \frac{n!}{(n-i)!i!} (-1)^{n-i} \mathcal{D}^i \left(\frac{d}{dx}\right)^{n-i} Y_-^0(u, x)v \tag{4.10}$$

for $u, v \in \mathbb{A} \oplus \mathbb{B}$, $n \in \mathbb{N}$.

The following proposition will play an important role for the rest of this section.

Proposition 4.2. For $a, a' \in \mathbb{A}$, $b, b' \in \mathbb{B}$, $n \in \mathbb{N}$, we have

$$Y_-^0(a, x) \mathcal{D}^n \otimes a' = 0 \quad (4.11)$$

$$Y_-^0(a, x) \mathcal{D}^n \otimes b = \sum_{i=0}^n \frac{n!}{(n-i)!} \mathcal{D}^{n-i} \otimes (a_0 b) x^{-1-i} \quad (4.12)$$

$$Y_-^0(b, x) \mathcal{D}^n \otimes a = \sum_{i=0}^n \frac{n!}{(n-i)!} \mathcal{D}^{n-i} \otimes (b_0 a) x^{-1-i} \quad (4.13)$$

$$\begin{aligned} Y_-^0(b, x) \mathcal{D}^n \otimes b' &= \sum_{i=0}^n \frac{n!}{(n-i)!} \mathcal{D}^{n-i} \otimes (b_0 b') x^{-1-i} \\ &\quad + \sum_{i=1}^{n+1} \frac{in!}{(n-i+1)!} \mathcal{D}^{n-i+1} \otimes (b_1 b') x^{-i-1} \end{aligned} \quad (4.14)$$

Proof. It follows immediately from (4.4)-(4.7), (4.10). $\square$

Next, we will show that $(C(\mathbb{A} \oplus \mathbb{B})/\hat{\partial}C(\mathbb{A}), Y_-^0, \mathcal{D})$ is a vertex Lie algebra.

Proposition 4.3. The subspace $\hat{\partial}C(\mathbb{A})$ of the nonassociative algebra $C(\mathbb{A} \oplus \mathbb{B})$ is a 2-sided ideal.

Proof. First we show that for $u \in \mathbb{A} \oplus \mathbb{B}$, $a \in \mathbb{A}$, $m, n \in \mathbb{N}$,

$$Y_-^0(\mathcal{D}^n \otimes u, x) \hat{\partial}(\mathcal{D}^m \otimes a) \in x^{-1} \hat{\partial}C(\mathbb{A})[x^{-1}].$$

Let $a, a' \in \mathbb{A}$, $m \in \mathbb{N}$, $n \geq 1$. By (3.1)-(3.2), (4.9), (4.11)-(4.13), we have

$$Y_-^0(a', x) \hat{\partial}(\mathcal{D}^m \otimes a) = Y_-^0(a', x) (\mathcal{D}^m \otimes \partial(a) - \mathcal{D}^{m+1} \otimes a) = 0,$$

and

$$\begin{aligned} &Y_-^0(\mathcal{D}^n \otimes a', x) \hat{\partial}(\mathcal{D}^m \otimes a) \\ &= Y_-^0(\mathcal{D}^n \otimes a', x) (\mathcal{D}^m \otimes \partial(a) - \mathcal{D}^{m+1} \otimes a) \\ &= \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^m Y_-^0(\partial(a), x) \mathcal{D}^n \otimes a') \\ &\quad - \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^{m+1} Y_-^0(a, -x) \mathcal{D}^n \otimes a') \\ &= 0. \end{aligned}$$

Let $a \in \mathbb{A}$, $b \in \mathbb{B}$, $m \in \mathbb{N}$, $n \geq 1$. By (3.1)-(3.2), (4.9), (4.12)-(4.14), we have

$$\begin{aligned}
& Y_-^0(b, x) \hat{\partial}(\mathcal{D}^m \otimes a) \\
&= Y_-^0(b, x) \mathcal{D}^m \otimes \partial(a) - Y_-^0(b, x) \mathcal{D}^{m+1} \otimes a \\
&= \sum_{i=0}^m \frac{m!}{(m-i)!} \mathcal{D}^{m-i} \otimes b_0 \partial(a) x^{-1-i} + \sum_{i=1}^{m+1} \frac{im!}{(m-i+1)!} \mathcal{D}^{m-i+1} \otimes b_1 \partial(a) x^{-i-1} \\
&\quad - \sum_{i=0}^{m+1} \frac{(m+1)!}{(m+1-i)!} \mathcal{D}^{m+1-i} \otimes b_0 a x^{-1-i} \\
&= \sum_{i=0}^m \frac{m!}{(m-i)!} \mathcal{D}^{m-i} \otimes \partial((\partial a)_1 b) x^{-1-i} + \sum_{i=1}^{m+1} \frac{im!}{(m-i+1)!} \mathcal{D}^{m-i+1} \otimes (\partial a)_1 b x^{-i-1} \\
&\quad - \sum_{i=0}^{m+1} \frac{(m+1)!}{(m+1-i)!} \mathcal{D}^{m+1-i} \otimes (\partial a)_1 b x^{-1-i} \\
&= \sum_{i=0}^m \frac{m!}{(m-i)!} \mathcal{D}^{m-i} \otimes \partial((\partial a)_1 b) x^{-1-i} - \sum_{i=0}^m \frac{m!}{(m-i)!} \mathcal{D}^{m+1-i} \otimes (\partial a)_1 b x^{-1-i} \\
&= \sum_{i=0}^m \frac{m!}{(m-i)!} \hat{\partial}(\mathcal{D}^{m-i} \otimes (\partial a)_1 b) x^{-1-i},
\end{aligned}$$

and

$$\begin{aligned}
& Y_-^0(\mathcal{D}^n \otimes b, x) \hat{\partial}(\mathcal{D}^m \otimes a) \\
&= Y_-^0(\mathcal{D}^n \otimes b, x) \mathcal{D}^m \otimes \partial(a) - Y_-^0(\mathcal{D}^n \otimes b, x) \mathcal{D}^{m+1} \otimes a \\
&= \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^m Y_-^0(\partial(a), -x) \mathcal{D}^n \otimes b) \\
&\quad - \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^{m+1} Y_-^0(a, -x) \mathcal{D}^n \otimes b) \\
&= \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^m \sum_{i=0}^n \frac{n!}{(n-i)!} \mathcal{D}^{n-i} \otimes (\partial(a)_0 b) (-x)^{-i-1}) \\
&\quad + \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^m \sum_{i=1}^{n+1} \frac{in!}{(n-i+1)!} \mathcal{D}^{n-i+1} \otimes \partial(a)_1 b (-x)^{-i-1}) \\
&\quad - \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^{m+1} \sum_{i=0}^n \frac{n!}{(n-i)!} \mathcal{D}^{n-i} \otimes (a_0 b) (-x)^{-1-i}) \\
&= 0.
\end{aligned}$$

These imply that for $u \in \mathbb{A} \oplus \mathbb{B}$, $a \in \mathbb{A}$, $m, n \in \mathbb{N}$,

$$Y_-^0(\mathcal{D}^n \otimes u, x) \hat{\partial}(\mathcal{D}^m \otimes a) \in x^{-1} \hat{\partial}C(\mathbb{A})[x^{-1}].$$

Next, we will show that for $u \in \mathbb{A} \oplus \mathbb{B}$, $a \in \mathbb{A}$, $m, n \in \mathbb{N}$,

$$Y_-^0(\hat{\partial}(\mathcal{D}^m \otimes a), x) \mathcal{D}^n \otimes u \in x^{-1} \hat{\partial}C(\mathbb{A})[x^{-1}].$$

Let $a, a' \in \mathbb{A}$, $b \in \mathbb{B}$, $n \in \mathbb{N}$. By (3.1), (4.9), (4.11), (4.13)-(4.14), we have

$$\begin{aligned}
Y_-^0(\hat{\partial}(a), x) \mathcal{D}^n \otimes a' &= Y_-^0(\partial(a), x) \mathcal{D}^n \otimes a' - Y(\mathcal{D} \otimes a, x) \mathcal{D}^n \otimes a' \\
&= -\text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n Y_-^0(a', -x) \mathcal{D} \otimes a) \\
&= 0,
\end{aligned}$$

and

$$\begin{aligned}
&Y_-^0(\hat{\partial}(1 \otimes a), x) \mathcal{D}^n \otimes b \\
&= Y_-^0(1 \otimes \partial a, x) \mathcal{D}^n \otimes b - Y_-^0(\mathcal{D} \otimes a, x) \mathcal{D}^n \otimes b \\
&= \sum_{i=0}^n \frac{n!}{(n-i)!} \mathcal{D}^{n-i} \otimes (\partial a)_{(0)} b x^{-1-i} + \sum_{i=1}^{n+1} \frac{in!}{(n-i+1)!} \mathcal{D}^{n-i+1} \otimes (\partial a)_1 b x^{-i-1} \\
&\quad - \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n Y_-^0(b, -x) \mathcal{D} \otimes a) \\
&= -\sum_{i=1}^{n+1} \frac{in!}{(n-i+1)!} \mathcal{D}^{n-i+1} \otimes a_0 b x^{-i-1} \\
&\quad + \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n (\mathcal{D} \otimes a_0 b (-x)^{-1} + 1 \otimes a_0 b (-x)^{-2})) \\
&= -\sum_{i=1}^{n+1} \frac{in!}{(n-i+1)!} \mathcal{D}^{n-i+1} \otimes a_0 b x^{-i-1} - \sum_{j=0}^n \frac{n!}{j!} \mathcal{D}^{j+1} \otimes a_0 b x^{-1-n+j} \\
&\quad + \sum_{j=0}^{n+1} \frac{(n+1)!}{j!} \mathcal{D}^j \otimes a_0 b x^{-2-n+j} \\
&= -\sum_{i=1}^{n+1} \frac{in!}{(n-i+1)!} \mathcal{D}^{n-i+1} \otimes a_0 b x^{-i-1} - \sum_{i=0}^n \frac{n!}{(n-i)!} \mathcal{D}^{n-i+1} \otimes a_0 b x^{-1-i} \\
&\quad + \sum_{i=0}^{n+1} \frac{(n+1)!}{(n-i+1)!} \mathcal{D}^{n-i+1} \otimes a_0 b x^{-1-i} \\
&= 0.
\end{aligned}$$

Let $a \in \mathbb{A}$, $b \in \mathbb{B}$, $n \in \mathbb{N}$, $m \geq 1$. By (3.1)-(3.2), (4.9), (4.11)-(4.12), we have

$$\begin{aligned}
Y_-^0(\hat{\partial}(\mathcal{D}^m \otimes a), x) \mathcal{D}^n \otimes a' &= Y_-^0(\mathcal{D}^m \otimes \partial a, x) \mathcal{D}^n \otimes a' - Y(\mathcal{D}^{m+1} \otimes a, x) \mathcal{D}^n \otimes a' \\
&= \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n Y_-^0(a', -x) \mathcal{D}^m \otimes \partial a) \\
&= 0.
\end{aligned}$$

Since $\mathcal{D}(\hat{\partial}(C(\mathbb{A}))) \subset \hat{\partial}(C(\mathbb{A}))$ and $Y_-^0(b, -x)\hat{\partial}(C(\mathbb{A})) \subset x^{-1}\hat{\partial}(C(\mathbb{A}))[x^{-1}]$, we have

$$\begin{aligned}
& Y_-^0(\hat{\partial}(\mathcal{D}^m \otimes a), x)\mathcal{D}^n \otimes b \\
&= Y_-^0(\mathcal{D}^m \otimes \partial(a), x)\mathcal{D}^n \otimes b - Y_-^0(\mathcal{D}^{m+1} \otimes a, x)\mathcal{D}^n \otimes b \\
&= \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n Y_-^0(b, -x)\mathcal{D}^m \otimes \partial(a)) \\
&\quad - \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n Y_-^0(b, -x)\mathcal{D}^{m+1} \otimes a) \\
&= \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n Y_-^0(b, -x)\hat{\partial}(\mathcal{D}^m \otimes a)) \in x^{-1}\hat{\partial}(C(\mathbb{A}))[x^{-1}].
\end{aligned}$$

It follows that for $u \in \mathbb{A} \oplus \mathbb{B}$, $a \in \mathbb{A}$, $m, n \in \mathbb{N}$, we have

$$Y_-^0(\hat{\partial}(\mathcal{D}^m \otimes a), x)\mathcal{D}^n \otimes u \in x^{-1}\hat{\partial}C(\mathbb{A})[x^{-1}].$$

Hence, $\hat{\partial}C(\mathbb{A})$ is a 2-sided ideal of $C(\mathbb{A} \oplus \mathbb{B})$. □

Set

$$\mathcal{C} = C(\mathbb{A} \oplus \mathbb{B})/\hat{\partial}C(\mathbb{A}). \quad (4.15)$$

Let ρ be a projection map:

$$C(\mathbb{A} \oplus \mathbb{B}) \rightarrow \mathcal{C}; \mathcal{D}^n \otimes u \mapsto \mathcal{D}^n \otimes u + \hat{\partial}C(\mathbb{A}).$$

For $u \in \mathbb{A} \oplus \mathbb{B}$, $n \in \mathbb{N}$, we set $\mathcal{D}^n u = \rho(\mathcal{D}^n \otimes u)$.

By (4.2), (4.3) and (4.15), we have the following.

Proposition 4.4. *For $a \in \mathbb{A}$, $n \in \mathbb{N}$, $\mathcal{D}^n(\partial a) = \mathcal{D}^{n+1}a$. Moreover, for $n \geq 1$, $\mathcal{C}_{(n)} = \{\mathcal{D}^{n-1}(b) \mid b \in \mathbb{B}\}$.*

Proposition 4.5. *For $u \in \mathbb{A} \oplus \mathbb{B}$, $i \geq 0$, $n \in \mathbb{N}$, $u_i \mathcal{C}_{(n)} \subset \mathcal{C}_{(n+\deg u-i-1)}$.*

Proof. This follows immediately from Propositions 4.2, 4.3. □

Next, we will show that $(\mathcal{C}, Y_-^0, \mathcal{D})$ is a vertex Lie algebra.

Lemma 4.6. *For $u, v \in \mathbb{A} \oplus \mathbb{B}$, $n \in \mathbb{N}$, we have*

$$Y_-^0(u, x)\mathcal{D}^n v = \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n Y_-^0(v, -x)u).$$

Proof. We first show that for $u, v \in \mathbb{A} \oplus \mathbb{B}$,

$$Y_-^0(u, x)v = \text{Sing}(e^{x\mathcal{D}} Y_-^0(v, -x)u).$$

Let $a, a' \in \mathbb{A}$, $b, b' \in \mathbb{B}$. By (3.2), (4.4)-(4.7), and Proposition 4.4, we have

$$\begin{aligned}
Y_-^0(a, x)a' &= 0 = \text{Sing}(e^{x\mathcal{D}} Y_-^0(a', -x)a), \\
Y_-^0(a, x)b &= a_0 b x^{-1} = -b_0 a x^{-1} = \text{Sing}(e^{x\mathcal{D}} Y_-^0(b, -x)a), \\
Y_-^0(b, x)a &= b_0 a x^{-1} = -a_0 b x^{-1} = \text{Sing}(e^{x\mathcal{D}} Y_-^0(a, -x)b),
\end{aligned}$$

and

$$\begin{aligned}
& \text{Sing}(e^{x\mathcal{D}}Y_-^0(b', -x)b) \\
&= \text{Sing}(e^{x\mathcal{D}}(-b'_0bx^{-1} + b'_1bx^{-2})) \\
&= -b'_0bx^{-1} + b'_1bx^{-2} + \mathcal{D}(b'_1b)x^{-1} \\
&= -b'_0bx^{-1} + b'_1bx^{-2} + \partial(b'_1b)x^{-1} \\
&= b_0b'x^{-1} + b_1b'x^{-2} \\
&= Y_-^0(b, x)b'.
\end{aligned}$$

These imply that for $u, v \in \mathbb{A} \oplus \mathbb{B}$,

$$Y_-^0(u, x)v = \text{Sing}(e^{x\mathcal{D}}Y_-^0(v, -x)u). \quad (4.16)$$

Next, we show that for $u, v \in \mathbb{A} \oplus \mathbb{B}$, $n \geq 1$,

$$Y_-^0(u, x)\mathcal{D}^n v = \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n Y_-^0(v, -x)u).$$

Clearly, for all $a, a' \in \mathbb{A}$, $n \geq 1$, we have

$$Y_-^0(a, x)\mathcal{D}^n a' = 0 = \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n Y_-^0(a', x)a).$$

Let $a \in \mathbb{A}$, $b \in \mathbb{B}$, $n \geq 1$. By (3.2), (4.6), (4.12), we have

$$\begin{aligned}
& \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n Y_-^0(b, -x)a) \\
&= \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n b_0a(-x)^{-1}) \\
&= -\sum_{i=0}^n \frac{n!}{i!} \mathcal{D}^i b_0a x^{-1-n+i} \\
&= -\sum_{i=0}^n \frac{n!}{(n-i)!} \mathcal{D}^{n-i} b_0a x^{-1-i} \\
&= Y_-^0(a, x)\mathcal{D}^n b.
\end{aligned}$$

Similarly, we have

$$Y_-^0(b, x)\mathcal{D}^n a = \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n Y_-^0(a, -x)b).$$

Let $b, b' \in \mathbb{B}$, $n \geq 1$. By (3.2), (4.7), (4.14), and Proposition 4.4, we have

$$\begin{aligned}
& \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n Y_-^0(b, -x)b') \\
&= \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n \{b_0 b'(-x)^{-1} + b_1 b'(-x)^{-2}\}) \\
&= \sum_{i=0}^n \frac{n!}{i!} \mathcal{D}^i (-b_0 b') x^{-1-n+i} + \sum_{i=0}^{n+1} \frac{(n+1)!}{i!} \mathcal{D}^i b_1 b' x^{-2-n+i} \\
&= \sum_{i=0}^n \frac{n!}{i!} \mathcal{D}^i (b'_0 b - \partial(b'_1 b)) x^{-1-n+i} + \sum_{i=0}^{n+1} \frac{(n+1)!}{i!} \mathcal{D}^i b_1 b' x^{-2-n+i} \\
&= \sum_{i=0}^n \frac{n!}{i!} (\mathcal{D}^i b'_0 b - \mathcal{D}^{i+1} b'_1 b) x^{-1-n+i} + \sum_{i=0}^{n+1} \frac{(n+1)!}{i!} \mathcal{D}^i b_1 b' x^{-2-n+i} \\
&= \sum_{j=0}^n \frac{n!}{(n-j)!} (\mathcal{D}^{n-j} b'_0 b - \mathcal{D}^{n-j+1} b'_1 b) x^{-1-j} + \sum_{j=0}^{n+1} \frac{(n+1)!}{(n+1-j)!} \mathcal{D}^{n+1-j} b_1 b' x^{-1-j} \\
&= Y_-^0(b', x) \mathcal{D}^n b.
\end{aligned}$$

Hence, we can conclude that $Y_-^0(u, x) \mathcal{D}^n v = \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^n Y_-^0(v, -x)u)$ for all $u, v \in \mathbb{A} \oplus \mathbb{B}$, $n \in \mathbb{N}$. $\square$

Proposition 4.7. For $u, v \in \mathbb{A} \oplus \mathbb{B}$, $n \in \mathbb{N}$,

$$[\mathcal{D}, Y_-^0(\mathcal{D}^n u, x)] e^{x_1 \mathcal{D}} v = \frac{d}{dx} Y_-^0(\mathcal{D}^n u, x) e^{x_1 \mathcal{D}} v = Y_-^0(\mathcal{D}(\mathcal{D}^n u), x) e^{x_1 \mathcal{D}} v. \quad (4.17)$$

Proof. We will follow the proof of Proposition 3.10 in [Li2] very closely. First we show that for $u, v \in \mathbb{A} \oplus \mathbb{B}$, $n \in \mathbb{N}$,

$$[\mathcal{D}, Y_-^0(\mathcal{D}^n u, x)] e^{x_1 \mathcal{D}} v = \frac{d}{dx} Y_-^0(\mathcal{D}^n u, x) e^{x_1 \mathcal{D}} v.$$

If we differentiate (4.8) with respect to x_1 , then we have

$$\begin{aligned}
Y_-^0(u, x) \mathcal{D} e^{x_1 \mathcal{D}} v &= \mathcal{D} e^{x_1 \mathcal{D}} e^{-x_1 \frac{d}{dx}} Y_-^0(u, x) v - e^{x_1 \mathcal{D}} \frac{d}{dx} e^{-x_1 \frac{d}{dx}} Y_-^0(u, x) v \\
&= \mathcal{D} Y_-^0(u, x) e^{x_1 \mathcal{D}} v - \frac{d}{dx} Y_-^0(u, x) e^{x_1 \mathcal{D}} v \quad (\text{by (4.8)}).
\end{aligned}$$

Hence, for $u, v \in \mathbb{A} \oplus \mathbb{B}$,

$$[\mathcal{D}, Y_-^0(u, x)] e^{x_1 \mathcal{D}} v = \frac{d}{dx} Y_-^0(u, x) e^{x_1 \mathcal{D}} v. \quad (4.18)$$

Let $u, v \in \mathbb{A} \oplus \mathbb{B}$, $m \in \mathbb{N}$, $n \geq 1$. By (2.1), (4.9), we have

$$\begin{aligned}
& [\mathcal{D}, Y_-^0(\mathcal{D}^n u, x)] \mathcal{D}^m v \\
&= \mathcal{D} Y_-^0(\mathcal{D}^n u, x) \mathcal{D}^m v - Y_-^0(\mathcal{D}^n u, x) \mathcal{D}^{m+1} v \\
&= \text{Sing}(\mathcal{D} e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^m Y_-^0(v, -x) \mathcal{D}^n u) - \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^{m+1} Y_-^0(v, -x) \mathcal{D}^n u) \\
&= \frac{d}{dx} \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^m Y_-^0(v, -x) \mathcal{D}^n u) \\
&= \frac{d}{dx} Y_-^0(\mathcal{D}^n u, x) \mathcal{D}^m v.
\end{aligned}$$

Hence, for $u, v \in \mathbb{A} \oplus \mathbb{B}$, $n \in \mathbb{N}$, $[\mathcal{D}, Y_-^0(\mathcal{D}^n u, x)] e^{x_1 \mathcal{D}} v = \frac{d}{dx} Y_-^0(\mathcal{D}^n u, x) e^{x_1 \mathcal{D}} v$.

Next, we will show that for $u, v \in \mathbb{A} \oplus \mathbb{B}$, $n \in \mathbb{N}$,

$$Y_-^0(\mathcal{D}(\mathcal{D}^n u), x) e^{x_1 \mathcal{D}} v = \frac{d}{dx} Y_-^0(\mathcal{D}^n u, x) e^{x_1 \mathcal{D}} v.$$

By (4.9), (4.18), and Lemma 4.6, we have that for $u, v \in \mathbb{A} \oplus \mathbb{B}$, $n, m \in \mathbb{N}$,

$$\begin{aligned}
Y_-^0(\mathcal{D}(\mathcal{D}^n u), x) \mathcal{D}^m v &= \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^m Y_-^0(v, -x) \mathcal{D}^{n+1} u) \\
&= \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^m (\mathcal{D} Y_-^0(v, -x) + \frac{d}{dx} Y_-^0(v, -x)) \mathcal{D}^n u) \\
&= \text{Sing}(e^{x\mathcal{D}} \mathcal{D} \left(-\frac{d}{dx}\right)^m Y_-^0(v, -x) \mathcal{D}^n u) \\
&\quad - \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^{m+1} Y_-^0(v, -x) \mathcal{D}^n u) \\
&= \frac{d}{dx} \text{Sing}(e^{x\mathcal{D}} \left(-\frac{d}{dx}\right)^m Y_-^0(v, -x) \mathcal{D}^n u) \\
&= \frac{d}{dx} Y_-^0(\mathcal{D}^n u, x) \mathcal{D}^m v.
\end{aligned}$$

Therefore, for $u, v \in \mathbb{A} \oplus \mathbb{B}$, $n \in \mathbb{N}$,

$$[\mathcal{D}, Y_-^0(\mathcal{D}^n u, x)] e^{x_1 \mathcal{D}} v = \frac{d}{dx} Y_-^0(\mathcal{D}^n u, x) e^{x_1 \mathcal{D}} v = Y_-^0(\mathcal{D}(\mathcal{D}^n u), x) e^{x_1 \mathcal{D}} v.$$

□

Corollary 4.8. For any $u \in \mathbb{A} \oplus \mathbb{B}$, $n \in \mathbb{N}$,

$$e^{x_1 \mathcal{D}} Y_-^0(\mathcal{D}^n u, x) e^{-x_1 \mathcal{D}} = e^{x_1 \frac{d}{dx}} Y_-^0(\mathcal{D}^n u, x) = Y_-^0(e^{x_1 \mathcal{D}} \mathcal{D}^n u, x) = Y_-^0(\mathcal{D}^n u, x + x_1).$$

Corollary 4.9. For $u \in \mathbb{A} \oplus \mathbb{B}$, $n \geq 1$,

$$(D^n u)_i = \begin{cases} 0, & \text{if } 0 \leq i < n; \\ (-1)^n \frac{i!}{(i-n)!} u_{i-n}, & \text{if } i \geq n. \end{cases}$$

Furthermore, we have $v_i \mathcal{C}_{(n)} \subset \mathcal{C}_{(n+m-i-1)}$ for all $v \in \mathcal{C}_{(m)}$, $i \in \mathbb{N}$.

Proof. It follows immediately from Proposition 4.5 and (4.17). $\square$

Corollary 4.10. For $u, v \in \mathbb{A} \oplus \mathbb{B}$, $n, m \in \mathbb{N}$,

$$Y_-^0(\mathcal{D}^n u, x) \mathcal{D}^m v = \text{Sing}(e^{x\mathcal{D}} Y_-^0(\mathcal{D}^m v, -x) \mathcal{D}^n u).$$

Proof. It follows from Lemma 4.6, (4.9), (4.17). $\square$

Proposition 4.11. For $u, v \in \mathbb{A} \oplus \mathbb{B}$, $m, n \in \mathbb{N}$,

$$[Y_-^0(\mathcal{D}^n u, x_1), Y_-^0(\mathcal{D}^m v, x_2)] = \text{Sing}(Y_-^0(Y_-^0(\mathcal{D}^n u, x_1 - x_2) \mathcal{D}^m v, x_2)).$$

Proof. We first show that for $u, v, w \in \mathbb{A} \oplus \mathbb{B}$,

$$\begin{aligned} & Y_-^0(u, x_1) Y_-^0(v, x_2) w - Y_-^0(v, x_2) Y_-^0(u, x_1) w \\ &= \text{Sing}(Y_-^0(Y_-^0(u, x_1 - x_2) v, x_2) w). \end{aligned}$$

By Proposition 2.6, it is enough to show that for $u, v, w \in \mathbb{A} \oplus \mathbb{B}$ from an ordered basis of $\mathbb{A} \oplus \mathbb{B}$ with $u \leq v \leq w$,

$$\begin{aligned} & Y_-^0(u, x_1) Y_-^0(v, x_2) w - Y_-^0(v, x_2) Y_-^0(u, x_1) w \\ &= \text{Sing}(Y_-^0(Y_-^0(u, x_1 - x_2) v, x_2) w). \end{aligned} \quad (4.19)$$

Let $\{a_i | i \in I\}$ and $\{b_j | j \in J\}$ be ordered bases of $\mathbb{A}$ and $\mathbb{B}$, respectively. We assume that $a_i \leq b_j$ for all $i \in I, j \in J$. Hence $C = \{a_i, b_j | i \in I, j \in J\}$ is an ordered basis of $\mathbb{A} \oplus \mathbb{B}$. Let $u \leq v \leq w \in C$. We claim that

$$\begin{aligned} & Y_-^0(u, x_1) Y_-^0(v, x_2) w - Y_-^0(v, x_2) Y_-^0(u, x_1) w \\ &= \text{Sing}(e^{x_2 \mathcal{D}} Y_-^0(w, -x_2) Y_-^0(u, x_1 - x_2) v). \end{aligned} \quad (4.20)$$

Case I: If $u, v, w \in \mathbb{A}$, by (4.4) all the three terms are straightly zero.

Case II: If $u, v \in \mathbb{A}$, $w \in \mathbb{B}$, by (4.4), (4.5) all the three terms are zero.

Case III: Assume that $u \in \mathbb{A}$, $v, w \in \mathbb{B}$. By (4.5)-(4.7), and (3.2)-(3.3), we have

$$\begin{aligned} & Y_-^0(u, x_1) Y_-^0(v, x_2) w - Y_-^0(v, x_2) Y_-^0(u, x_1) w \\ &= Y_-^0(u, x_1) (v_0 w x_2^{-1} + v_1 w x_2^{-2}) - Y_-^0(v, x_2) (u_0 w x_1^{-1}) \\ &= u_0 v_0 w x_1^{-1} x_2^{-1} - v_0 u_0 w x_1^{-1} x_2^{-1} \\ &= (u_0 v)_0 w x_1^{-1} x_2^{-1}, \end{aligned}$$

and

$$\begin{aligned} & \text{Sing}(e^{x_2 \mathcal{D}} Y_-^0(w, -x_2) Y_-^0(u, x_1 - x_2) v) \\ &= \text{Sing}(e^{x_2 \mathcal{D}} w_0 u_0 v (-x_2)^{-1} (x_1 - x_2)^{-1}) \\ &= -w_0 u_0 v x_1^{-1} x_2^{-1} \\ &= (u_0 v)_0 w x_1^{-1} x_2^{-1}. \end{aligned}$$

Hence, (4.20) holds when $u \in \mathbb{A}, v, w \in \mathbb{B}$.

Case IV: Assume that $u, v, w \in \mathbb{B}$. By (3.1)-(3.3), (4.6)-(4.7), and Proposition 4.4, we have

$$\begin{aligned}
& Y_-^0(u, x_1)Y_-^0(v, x_2)w - Y_-^0(v, x_2)Y_-^0(u, x_1)w \\
&= Y_-^0(u, x_1)(v_0wx_2^{-1} + v_1wx_2^{-2}) - Y_-^0(v, x_2)(u_0wx_1^{-1} + u_1wx_1^{-2}) \\
&= u_0v_0wx_1^{-1}x_2^{-1} + u_1v_0wx_1^{-2}x_2^{-1} + u_0v_1wx_1^{-1}x_2^{-2} \\
&\quad - v_0u_0wx_1^{-1}x_2^{-1} - v_1u_0wx_1^{-1}x_2^{-2} - v_0u_1wx_1^{-2}x_2^{-1} \\
&= (u_0v)_0wx_1^{-1}x_2^{-1} - (v_0u)_1wx_1^{-2}x_2^{-1} + (u_0v)_1wx_1^{-1}x_2^{-2},
\end{aligned}$$

and

$$\begin{aligned}
& \text{Sing}(e^{x_2\mathcal{D}}Y_-^0(w, -x_2)Y_-^0(u, x_1 - x_2)v) \\
&= \text{Sing}(e^{x_2\mathcal{D}}(w_0u_0v(-x_2)^{-1}(x_1 - x_2)^{-1} + w_1u_0v(-x_2)^{-2}(x_1 - x_2)^{-1})) \\
&\quad + \text{Sing}(e^{x_2\mathcal{D}}w_0u_1v(-x_2)^{-1}(x_1 - x_2)^{-2}) \\
&= -w_0u_0vx_1^{-1}x_2^{-1} + w_1u_0vx_1^{-1}x_2^{-2} + w_1u_0vx_1^{-2}x_2^{-1} + \partial(w_1u_0v)x_1^{-1}x_2^{-1} \\
&\quad - w_0u_1vx_1^{-2}x_2^{-1} \\
&= (u_0v)_0wx_1^{-1}x_2^{-1} + (u_0v)_1wx_1^{-1}x_2^{-2} + ((u_0v)_1w - w_0u_1v)x_1^{-2}x_2^{-1} \\
&= (u_0v)_0wx_1^{-1}x_2^{-1} + (u_0v)_1wx_1^{-1}x_2^{-2} + (-(v_0u)_1w + (\partial(v_1u))_1w + (u_1v)_0w)x_1^{-2}x_2^{-1} \\
&= (u_0v)_0wx_1^{-1}x_2^{-1} + (u_0v)_1wx_1^{-1}x_2^{-2} + (-(v_0u)_1w - (v_1u)_0w + (u_1v)_0w)x_1^{-2}x_2^{-1} \\
&= (u_0v)_0wx_1^{-1}x_2^{-1} + (u_0v)_1wx_1^{-1}x_2^{-2} - (v_0u)_1wx_1^{-2}x_2^{-1}.
\end{aligned}$$

Hence, (4.20) holds when $u, v, w \in \mathbb{B}$.

Therefore, for $u, v, w \in C$ such that $u \leq v \leq w$, we have

$$Y_-^0(u, x_1)Y_-^0(v, x_2)w - Y_-^0(v, x_2)Y_-^0(u, x_1)w = \text{Sing}(e^{x_2\mathcal{D}}Y_-^0(w, -x_2)Y_-^0(u, x_1 - x_2)v).$$

By Corollary 4.10 and Proposition 2.1, we also have

$$\begin{aligned}
[Y_-^0(u, x_1), Y_-^0(v, x_2)]w &= \text{Sing}(e^{x_2\mathcal{D}}\text{Sing}(e^{-x_2\mathcal{D}}Y_-^0(Y_-^0(u, x_1 - x_2)v, x_2)w)) \\
&= \text{Sing}(Y_-^0(Y_-^0(u, x_1 - x_2)v, x_2)w)
\end{aligned} \tag{4.21}$$

for all $u, v, w \in C$ such that $u \leq v \leq w$. Moreover, (4.21) holds for all $u, v \in \mathbb{A} \oplus \mathbb{B}$.

Next, we will show that for $u, v \in \mathbb{A} \oplus \mathbb{B}$, $m, n \in \mathbb{N}$,

$$[Y_-^0(\mathcal{D}^n u, x_1), Y_-^0(\mathcal{D}^m v, x_2)] = \text{Sing}(Y_-^0(Y_-^0(\mathcal{D}^n u, x_1 - x_2)\mathcal{D}^m v, x_2)).$$

We will follow the last part of the proof of Theorem 3.6 in [Li2]. Let $u, v \in \mathbb{A} \oplus \mathbb{B}$.

By Corollary 4.8, we have

$$\begin{aligned}
& [Y_-^0(e^{z_1\mathcal{D}}u, x_1), Y_-^0(e^{z_2\mathcal{D}}v, x_2)]e^{z\mathcal{D}}w \\
&= e^{z_1\frac{\partial}{\partial x_1}}e^{z_2\frac{\partial}{\partial x_2}}[Y_-^0(u, x_1), Y_-^0(v, x_2)]e^{z\mathcal{D}}w \\
&= e^{(z_1-z)\frac{\partial}{\partial x_1}}e^{(z_2-z)\frac{\partial}{\partial x_2}}e^{z\mathcal{D}}[Y_-^0(u, x_1), Y_-^0(v, x_2)]w \\
&= e^{(z_1-z)\frac{\partial}{\partial x_1}}e^{(z_2-z)\frac{\partial}{\partial x_2}}e^{z\mathcal{D}}\text{Sing}(Y_-^0(Y_-^0(u, x_1-x_2)v, x_2)w) \\
&= e^{(z_1-z)\frac{\partial}{\partial x_1}}e^{(z_2-z)\frac{\partial}{\partial x_2}}\text{Sing}_{x_1, x_2}(Y_-^0(Y_-^0(u, x_1-x_2)v, x_2+z)e^{z\mathcal{D}}w) \\
&= \text{Sing}_{x_1, x_2}(Y_-^0(Y_-^0(u, x_1-x_2+z_1-z_2)v, x_2+z_2)e^{z\mathcal{D}}w) \\
&= \text{Sing}_{x_1, x_2}(Y_-^0(Y_-^0(e^{(z_1-z_2)\mathcal{D}}u, x_1-x_2)v, x_2+z_2)e^{z\mathcal{D}}w) \\
&= \text{Sing}_{x_1, x_2}(Y_-^0(e^{-z_2\mathcal{D}}Y_-^0(e^{z_1\mathcal{D}}u, x_1-x_2)e^{z_2\mathcal{D}}v, x_2+z_2)e^{z\mathcal{D}}w) \\
&= \text{Sing}_{x_1, x_2}Y_-^0(Y_-^0(e^{z_1\mathcal{D}}u, x_1-x_2)e^{z_2\mathcal{D}}v, x_2)e^{z\mathcal{D}}w).
\end{aligned}$$

Hence, $[Y_-^0(\mathcal{D}^n u, x_1), Y_-^0(\mathcal{D}^m v, x_2)] = \text{Sing}(Y_-^0(Y_-^0(\mathcal{D}^n u, x_1-x_2)\mathcal{D}^m v, x_2))$. $\square$

Theorem 4.12. $(\mathcal{C}, Y_-^0, \mathcal{D})$ is a vertex Lie algebra. Moreover, $S(\mathcal{C})$ is a vertex Poisson algebra. In particular, Y_-^0 extend to a vertex Poisson algebra structure Y_- on $S(\mathcal{C})$ in the following way. First for $u \in \mathcal{C}$, we define a unique element

$$\widetilde{Y_-^0}(u, x) \in x^{-1}(\text{Der } S(\mathcal{C}))[[x^{-1}]]$$

by

$$\widetilde{Y_-^0}(u, x)v = Y_-^0(u, x)v \quad \text{for } u, v \in \mathcal{C}.$$

For $a \in S(\mathcal{C})$, we define

$$Y_-(a, x) \in x^{-1}(\text{Der } S(\mathcal{C}))[[x^{-1}]]$$

by

$$Y_-(a, x)u = \text{Sing}(e^{x\mathcal{D}}\widetilde{Y_-^0}(u, -x)a) \quad \text{for } u \in \mathcal{C}.$$

Proof. It follows from Proposition 4.7, Corollary 4.10, Proposition 4.11, Proposition 2.16, and Proposition 3.7 in [Li2]. $\square$

Corollary 4.13. Let $u \in \mathcal{C}$. We have $Y_-(u, x)v = Y_-^0(u, x)v$ for all $v \in \mathcal{C}$. Furthermore, $Y_-(u, x)a = \widetilde{Y_-^0}(u, x)a$ for all $a \in S(\mathcal{C})$.

Since $\mathcal{C}$ is an $\mathbb{N}$ -graded vector space, it implies that $S(\mathcal{C}) = \coprod_{n \in \mathbb{N}} S(\mathcal{C})_{(n)}$ is an $\mathbb{N}$ -graded vector space. By Corollary 4.9, Theorem 4.12 and Corollary 4.13, we can conclude that:

Lemma 4.14. For $u \in \mathcal{C}_{(m)}$, $i \in \mathbb{N}$, $u_i S(\mathcal{C})_{(n)} \subset S(\mathcal{C})_{(n+m-i-1)}$. Here $u_i \in \text{Der } S(\mathcal{C})$.

Theorem 4.15. $(S(\mathcal{C}), \mathcal{D})$ is an $\mathbb{N}$ -graded vertex Poisson algebra. In particular, we have $S(\mathcal{C})_{(0)}$ is $S(\mathbb{A})$ and for $n \geq 1$,

$$\begin{aligned} S(\mathcal{C})_{(n)} &= \text{span}_{\mathbb{C}}\{ \mathcal{D}^{n_1}(b_1) \cdot \mathcal{D}^{n_2}(b_2) \cdot \dots \cdot \mathcal{D}^{n_k}(b_k) \cdot a_1 \cdot \dots \cdot a_l \mid a_j \in \mathbb{A}, b_i \in \mathbb{B}, \\ &\quad l \in \mathbb{N}, k \geq 1, n_1 \geq \dots \geq n_k \geq 0, n_1 + \dots + n_k + k = n \}. \end{aligned}$$

Here, $S(\mathbb{A})$ is the symmetric algebra over the space $\mathbb{A}$.

Proof. First, we show that $\mathcal{D}S(\mathcal{C})_{(n)} \subset S(\mathcal{C})_{(n+1)}$ for $n \in \mathbb{N}$. Clearly, for $n \in \mathbb{N}$, $\mathcal{D}\mathcal{C}_{(n)} \subset \mathcal{C}_{(n+1)}$. Since $\mathcal{D}$ is a derivation on $S(\mathcal{C})$, we can conclude immediately that $\mathcal{D}S(\mathcal{C})_{(n)} \subset S(\mathcal{C})_{(n+1)}$.

Next, let $a \in S(\mathcal{C})_{(n)}$, $i \in \mathbb{N}$. We will show that

$$a_i S(\mathcal{C})_{(r)} \subset S(\mathcal{C})_{(r+n-i-1)} \quad \text{for all } r \in \mathbb{N}.$$

Let $r \in \mathbb{N}$ and let $u \in \mathcal{C}_{(r)}$. By Corollary 4.13 and Lemma 4.14, we have

$$\begin{aligned} &Y_-(a, x)u \\ &= \text{Sing}(e^{x\mathcal{D}} \widetilde{Y}_0^-(u, -x)a) \\ &= \text{Sing}(e^{x\mathcal{D}} \sum_{j=0}^{\deg a + \deg u - 1} u_j a(-x)^{-j-1}) \\ &= \sum_{j=0}^{\deg a + \deg u - 1} \sum_{k=0}^j (-1)^{-j-1} \frac{\mathcal{D}^{j-k}}{(j-k)!} (u_j a) x^{-1-k}. \end{aligned}$$

Here, $u_j \in \text{Der } S(\mathcal{C})$. It implies that $a_i \mathcal{C}_{(r)} \subset S(\mathcal{C})_{(r+n-i-1)}$. Since a_i is a derivation on $S(\mathcal{C})$ and $a_i \mathcal{C}_{(m)} \subset S(\mathcal{C})_{(m+n-i-1)}$ for all $m \in \mathbb{N}$, it follows that $a_i S(\mathcal{C})_{(m)} \subset S(\mathcal{C})_{(m+n-i-1)}$ for all $m \in \mathbb{N}$. Therefore, $(S(\mathcal{C}), Y_-)$ is an $\mathbb{N}$ -graded vertex Poisson algebra.

The second statement is clear. $\square$

5 Vertex Poisson algebras associated with Courant algebroids

In this section, we construct an $\mathbb{N}$ -graded vertex Poisson algebra $S(\mathcal{C})_{\mathcal{B}} = \coprod_{n \in \mathbb{N}} (S(\mathcal{C})_{\mathcal{B}})_{(n)}$ associated with a Courant $\mathcal{A}$ -algebroid $\mathcal{B}$. Also, we show that $(S(\mathcal{C})_{\mathcal{B}})_{(0)}$ can be naturally identified with $\mathcal{A}$ as a commutative associative algebra and $(S(\mathcal{C})_{\mathcal{B}})_{(1)}$ can be identified with $\mathcal{B}$ as a Courant $\mathcal{A}$ -algebroid.

For the rest of this paper, we assume that $\mathcal{A}$ is a commutative associative algebra with the identity e and $\mathcal{B}$ is a Courant $\mathcal{A}$ -algebroid. By Proposition 3.6, $\mathcal{A} \oplus \mathcal{B}$ is a 1-truncated conformal algebra such that for $a, a' \in \mathcal{A}$, $u, v \in \mathcal{B}$,

$$(au)_0 a' = a(u_0 a'), \quad (5.1)$$

$$(au)_1 v = a(u_1 v) = u_1(av), \quad (5.2)$$

$$u_0(av) = a(u_0 v) + (u_0 a)v, \quad (5.3)$$

$$u_0(aa') = a(u_0 a') + (u_0 a)a'. \quad (5.4)$$

Furthermore, by Theorem 4.15, we have an $\mathbb{N}$ -graded vertex Poisson algebra $S(\mathcal{C}) = \coprod_{n \in \mathbb{N}} S(\mathcal{C})_{(n)}$ associated with a 1-truncated conformal algebra $\mathcal{A} \oplus \mathcal{B}$. We denote a multiplication on $S(\mathcal{C})$ by " $\cdot$ ". Let

$$E_0 = \text{span}\{ e - 1, \ a \cdot a' - aa' \mid a, \ a' \in \mathcal{A} \} \subset S(\mathcal{C})_{(0)}, \quad (5.5)$$

$$E_1 = \text{span}\{ a \cdot b - ab \mid a \in \mathcal{A}, \ b \in \mathcal{B} \} \subset S(\mathcal{C})_{(1)}, \quad (5.6)$$

$$E = E_0 \oplus E_1 \subset S(\mathcal{C}). \quad (5.7)$$

Lemma 5.1. *For $u \in \mathcal{A} \oplus \mathcal{B}$, $n \in \mathbb{N}$, we have $u_n E \subset E$. Moreover, $\mathcal{D}E_0 \subset E_1$.*

Proof. We first show that for $u \in \mathcal{A} \oplus \mathcal{B}$, $n \in \mathbb{N}$,

$$u_n E_0 \subset E_0.$$

For $a \in \mathcal{A}$, $n \in \mathbb{N}$, we have $a_n E_0 = 0$. Similarly, for $b \in \mathcal{B}$, $n \geq 1$, we have $b_n E_0 = 0$. Let $b \in \mathcal{B}$. By Corollary 2.9, Corollary 3.7, and (5.4), we have

$$b_0(e - 1) = 0$$

and

$$\begin{aligned} & b_0(a \cdot a' - aa') \\ &= a \cdot b_0(a') + b_0(a) \cdot a' - (b_0 a) a' - a(b_0 a') \\ &= (a \cdot (b_0 a') - a(b_0 a')) + ((b_0 a) \cdot a' - (b_0 a) a') \in E_0. \end{aligned}$$

for all $a, a' \in \mathcal{A}$. Therefore, for $u \in \mathcal{A} \oplus \mathcal{B}$, $n \in \mathbb{N}$, $u_n E_0 \subset E_0$.

Next, we show that for $u \in \mathcal{A} \oplus \mathcal{B}$, $n \in \mathbb{N}$, $u_n E_1 \subset E$. Clearly, for $a \in \mathcal{A}$, $b \in \mathcal{B}$, $m \geq 1$, $n \geq 2$, we have

$$a_m E_1 \subset E_1, \text{ and } b_n E_1 \subset E_1.$$

Let $a', a \in \mathcal{A}$, $b \in \mathcal{B}$. By (3.2), (5.1), we have

$$\begin{aligned} & a'_0(a \cdot b - ab) \\ &= a'_0(a) \cdot b + a \cdot a'_0(b) + (ab)_0 a' \\ &= a \cdot a'_0(b) + a(b_0 a') \\ &= a \cdot (a'_0 b) - a(a'_0 b) \in E_0. \end{aligned}$$

Hence,

$$a_n E_1 \subset E \quad \text{for all } a \in \mathcal{A}, \ n \in \mathbb{N}.$$

Let $a \in \mathcal{A}$, $u, b \in \mathcal{B}$. By (5.2), (5.3), we have

$$u_0(a \cdot b - ab) = (u_0 a) \cdot b + a \cdot (u_0 b) - a(u_0 b) - (u_0 a)b \in E_1,$$

and

$$u_1(a \cdot b - ab) = (u_1 a) \cdot b + a \cdot (u_1 b) - u_1(ab) = a \cdot u_1(b) - a(u_1 b) \in E_0.$$

It follows that for $b \in \mathcal{B}$, $n \in \mathbb{N}$, $b_n E_1 \subset E$.

Next, we show that $\mathcal{D}E_0 \subset E_1$. By Proposition 4.4, we have that

$$\mathcal{D}(e - 1) = \mathcal{D}e = \partial(e) = 0 \in E_1.$$

Let $a, a' \in A$. By Proposition 4.4, and the fact that $\mathcal{D}$ is a derivation on $S(\mathcal{C})$ and ∂ is a derivation from $\mathcal{A}$ to $\mathcal{B}$, we have

$$\begin{aligned} & \mathcal{D}(a \cdot a' - aa') \\ = & a \cdot (\mathcal{D}a') + (\mathcal{D}a) \cdot a' - \mathcal{D}(aa') \\ = & a \cdot \partial(a') + \partial(a) \cdot a' - \partial(aa') \\ = & a \cdot \partial(a') + a' \cdot \partial(a) - a\partial(a') - a'\partial(a) \in E_1. \end{aligned}$$

Therefore, $\mathcal{D}E_0 \subset E_1$. □

For a subset U of $S(\mathcal{C})$, we set

$$\mathbb{C}[\mathcal{D}](U) = \text{span}\{ \mathcal{D}^m(u) \mid u \in U, m \in \mathbb{N} \}.$$

Define

$$I_{\mathcal{B}} = S(\mathcal{C}) \cdot \mathbb{C}[\mathcal{D}](E) \subset S(\mathcal{C}),$$

an ideal of a commutative associative algebra $S(\mathcal{C})$.

Lemma 5.2. *$I_{\mathcal{B}}$ is an ideal of a vertex Poisson algebra $S(\mathcal{C})$. Moreover,*

$$\begin{aligned} S(\mathcal{C})_{(0)} &= (I_{\mathcal{B}})_{(0)} \oplus \mathcal{A} \text{ and,} \\ S(\mathcal{C})_{(1)} &= (I_{\mathcal{B}})_{(1)} \oplus \mathcal{B}. \end{aligned}$$

Proof. Since $\mathcal{D}$ is a derivation on $S(\mathcal{C})$ and $\mathcal{D}(\mathbb{C}[\mathcal{D}](E)) \subset \mathbb{C}[\mathcal{D}](E)$, it follows that

$$\mathcal{D}I_{\mathcal{B}} \subset I_{\mathcal{B}}. \tag{5.8}$$

Let $v \in \mathcal{A} \oplus \mathcal{B}$, and $u \in E$. We claim that for $m \in \mathbb{N}$,

$$v_i \mathcal{D}^m(u) \in \mathbb{C}[\mathcal{D}](E) \quad \text{for all } i \in \mathbb{N}.$$

We prove this by an induction on m . For the case when $m = 0$, it follows immediately from Lemma 5.1. Recall that for $i \in \mathbb{N}$, $[\mathcal{D}, v_i] = -iv_{i-1}$ (see Proposition 2.5). We now assume that for $n < m$, $v_i \mathcal{D}^n(u) \in \mathbb{C}[\mathcal{D}](E)$ for all $i \in \mathbb{N}$. Observe that

$$v_i \mathcal{D}^m(u) = \mathcal{D}v_i \mathcal{D}^{m-1}(u) + iv_{i-1} \mathcal{D}^{m-1}(u).$$

By an induction hypothesis, we conclude that $v_i \mathcal{D}^m(u) \in \mathbb{C}[\mathcal{D}](E)$ for all $i \in \mathbb{N}$. Therefore, for $m \in \mathbb{N}$,

$$v_i \mathcal{D}^m(u) \in \mathbb{C}[\mathcal{D}](E) \text{ for all } i \in \mathbb{N}. \tag{5.9}$$

Next, we show that for $v \in \mathcal{A} \oplus \mathcal{B}$, $i \in \mathbb{N}$, $v_i I_{\mathcal{B}} \subset I_{\mathcal{B}}$. Let $v \in \mathcal{A} \oplus \mathcal{B}$ and $i \in \mathbb{N}$. By (2.12), (5.9), we have that for $a \in S(\mathcal{C})$, $u \in E$, $m \in \mathbb{N}$,

$$v_i(a \cdot \mathcal{D}^m u) = a \cdot v_i(\mathcal{D}^m u) + v_i(a) \cdot \mathcal{D}^m u \in I_{\mathcal{B}}.$$

This implies that for $v \in \mathcal{A} \oplus \mathcal{B}$, $i \in \mathbb{N}$,

$$v_i I_{\mathcal{B}} \subset I_{\mathcal{B}}. \quad (5.10)$$

Next, we will show that for $a \in S(\mathcal{C})$, $w \in I_{\mathcal{B}}$, $i \in \mathbb{N}$,

$$a_i w \in I_{\mathcal{B}}.$$

By (2.8), (5.8), (5.10), we have that for $w \in I_{\mathcal{B}}$, $v \in \mathcal{A} \oplus \mathcal{B}$, $i \in \mathbb{N}$,

$$w_i v \in I_{\mathcal{B}}.$$

Furthermore, by (5.10), Corollary 4.9, (2.8), and (5.8), we conclude that

$$(\mathcal{D}^m v)_i w \in I_{\mathcal{B}}, \quad \text{and } w_i (\mathcal{D}^m v) \in I_{\mathcal{B}}$$

for all $v \in \mathcal{A} \oplus \mathcal{B}$, $w \in I_{\mathcal{B}}$, $i \in \mathbb{N}$, and $m \geq 1$. Hence, for $w \in I_{\mathcal{B}}$, $i \in \mathbb{N}$,

$$w_i \mathcal{C} \subset I_{\mathcal{B}}.$$

Let $w \in I_{\mathcal{B}}$, $i \in \mathbb{N}$. Since w_i is a derivation on $S(\mathcal{C})$ and $w_i \mathcal{C} \subset I_{\mathcal{B}}$, it follows that

$$w_i S(\mathcal{C}) \subset I_{\mathcal{B}}. \quad (5.11)$$

Hence, by (2.8), (5.8), (5.11), we can conclude that for $a \in S(\mathcal{C})$, $w \in I_{\mathcal{B}}$, $i \in \mathbb{N}$, $a_i w \in I_{\mathcal{B}}$ and $I_{\mathcal{B}}$ is an ideal of $S(\mathcal{C})$ as a vertex Poisson algebra.

Next, we will show that $S(\mathcal{C})_{(0)} = (I_{\mathcal{B}})_{(0)} \oplus \mathcal{A}$ and $S(\mathcal{C})_{(1)} = (I_{\mathcal{B}})_{(1)} \oplus \mathcal{B}$. Clearly,

$$(I_{\mathcal{B}})_{(0)} = S(\mathcal{A})E_0 \text{ and } (I_{\mathcal{B}})_{(1)} = S(\mathcal{A})\mathcal{D}E_0 + S(\mathcal{A}) \cdot \mathcal{B} \cdot E_0 + S(\mathcal{A}) \cdot E_1.$$

By Lemma 5.1, we have $S(\mathcal{A})\mathcal{D}E_0 \subset S(\mathcal{A}) \cdot E_1$. Since $\mathcal{B}$ is an $\mathcal{A}$ -module, it implies that for $a, a' \in \mathcal{A}$, $b \in \mathcal{B}$

$$\begin{aligned} & b \cdot (a \cdot a') - b \cdot (aa') \\ = & (a \cdot a') \cdot b - (aa') \cdot b \\ = & a \cdot (a' \cdot b) - a \cdot (a'b) + a \cdot (a'b) - (aa') \cdot b \\ = & a \cdot (a' \cdot b) - a \cdot (a'b) + a \cdot (a'b) - (aa') \cdot b + (aa')b - (aa')b \\ = & a \cdot (a' \cdot b) - a \cdot (a'b) + a \cdot (a'b) - (aa') \cdot b + (aa')b - a(a'b) \in S(\mathcal{A}) \cdot E_1. \end{aligned}$$

Therefore,

$$S(\mathcal{A}) \cdot \mathcal{B} \cdot E_0 \subset S(\mathcal{A}) \cdot E_1, \text{ and } (I_{\mathcal{B}})_{(1)} = S(\mathcal{A}) \cdot E_1.$$

Moreover, we have

$$S(\mathcal{C})_{(0)} = S(\mathcal{A}) = \mathcal{A} \oplus (I_{\mathcal{B}})_{(0)},$$

and

$$S(\mathcal{C})_{(1)} = S(\mathcal{A}) \cdot \mathcal{B} = \mathcal{B} \oplus (I_{\mathcal{B}})_{(1)}.$$

□

Set

$$S(\mathcal{C})_{\mathcal{B}} = S(\mathcal{C})/I_{\mathcal{B}},$$

an $\mathbb{N}$ -graded vertex Poisson algebra.

Theorem 5.3. *Let $\mathcal{B}$ be a Courant $\mathcal{A}$ -algebroid and let $S(\mathcal{C})_{\mathcal{B}} = \coprod_{n \in \mathbb{N}} (S(\mathcal{C})_{\mathcal{B}})_{(n)}$ be the associated $\mathbb{N}$ -graded vertex Poisson algebra. We have that $(S(\mathcal{C})_{\mathcal{B}})_{(0)} = \mathcal{A}$ and $(S(\mathcal{C})_{\mathcal{B}})_{(1)} = \mathcal{B}$. Moreover, for any $n \geq 1$,*

$$(S(\mathcal{C})_{\mathcal{B}})_{(n)} = \text{span}\{ \mathcal{D}^{n_1}(b_1) \cdot \mathcal{D}^{n_2}(b_2) \cdot \dots \cdot \mathcal{D}^{n_k}(b_k) \mid b_i \in \mathcal{B}, n_1 \geq \dots \geq n_k \geq 0, k \geq 1, n_1 + \dots + n_k + k = n \}.$$

Proof. By Lemma 5.2, we have $(S(\mathcal{C})_{\mathcal{B}})_{(0)} = \mathcal{A}$ and $(S(\mathcal{C})_{\mathcal{B}})_{(1)} = \mathcal{B}$. For a subset U of $S(\mathcal{C})_{\mathcal{B}}$, we set

$$\mathbb{C}[\mathcal{D}]U = \text{Span}\{ \mathcal{D}^m(u) \mid u \in U, m \in \mathbb{N} \}.$$

We will show that

$$\coprod_{n \geq 1} (S(\mathcal{C})_{\mathcal{B}})_{(n)} = S(\mathbb{C}[\mathcal{D}]\mathcal{B}) \cdot \mathbb{C}[\mathcal{D}]\mathcal{B}.$$

Here, $S(\mathbb{C}[\mathcal{D}]\mathcal{B})$ is a symmetric algebra over the space $\mathbb{C}[\mathcal{D}]\mathcal{B}$. By Theorem 4.15, it is enough to show that $(\mathbb{C}[\mathcal{D}]\mathcal{B}) \cdot \mathcal{A} \subset \mathbb{C}[\mathcal{D}]\mathcal{B}$. Clearly, for $a \in \mathcal{A}$, $b \in \mathcal{B}$, we have

$$b \cdot a = a \cdot b = ab + a \cdot b - ab = ab \in \mathcal{B}.$$

Recall that for $u, v \in S(\mathcal{C})_{\mathcal{B}}$, $n \in \mathbb{N}$, we have

$$\mathcal{D}^n(u \cdot v) = \sum_{i=0}^n \binom{n}{i} \mathcal{D}^i(u) \cdot \mathcal{D}^{n-i}(v).$$

Let $a \in \mathcal{A}$, $b \in \mathcal{B}$, $n \geq 1$. By Proposition 4.4, we have

$$\begin{aligned} (\mathcal{D}^n b) \cdot a &= \mathcal{D}^n(a \cdot b) - \sum_{i=1}^n \binom{n}{i} \mathcal{D}^i(a) \cdot \mathcal{D}^{n-i}(b) \\ &= \mathcal{D}^n(ab) - \sum_{i=1}^n \binom{n}{i} \mathcal{D}^{i-1}(\partial(a)) \cdot \mathcal{D}^{n-i}(b). \end{aligned}$$

Therefore, for $a \in \mathcal{A}$, $b \in \mathcal{B}$ and $n \in \mathbb{N}$, we have $(\mathcal{D}^n b) \cdot a \in S(\mathbb{C}[\mathcal{D}]\mathcal{B}) \cdot \mathbb{C}[\mathcal{D}]\mathcal{B}$. It follows that

$$\coprod_{n \geq 1} (S(\mathcal{C})_{\mathcal{B}})_{(n)} = S(\mathbb{C}[\mathcal{D}]\mathcal{B}) \cdot \mathbb{C}[\mathcal{D}]\mathcal{B}.$$

□

6 Appendix

Definition 6.1. [LLi] A vertex algebra is a vector space V equipped with a linear map

$$\begin{aligned} Y : V &\rightarrow (\text{End } V)[[x, x^{-1}]], \\ v &\mapsto Y(v, x) = \sum_{n \in \mathbb{Z}} v_n x^{-n-1} \quad (\text{where } v_n \in \text{End } V) \end{aligned} \quad (6.1)$$

and equipped with a distinguished vector $\mathbf{1}$, called the vacuum (vector), such that for $u, v \in V$,

$$u_n v = 0 \quad \text{for } n \text{ sufficiently large,} \quad (6.2)$$

$$Y(\mathbf{1}, x) = 1, \quad (6.3)$$

$$Y(v, x)\mathbf{1} \in V[[x]] \quad \text{and} \quad \lim_{x \rightarrow 0} Y(v, x)\mathbf{1} = v \quad (6.4)$$

and

$$\begin{aligned} &x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) Y(u, x_1) Y(v, x_2) - x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) Y(v, x_2) Y(u, x_1) \\ &= x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) Y(Y(u, x_0)v, x_2) \end{aligned} \quad (6.5)$$

the Jacobi identity.

References

- [BeD] A. Beilinson, and V. Drinfeld, Chiral Algebras, Colloquium Publications, **Vol. 51**, Amer. Math. Soc., Providence, 2004.
- [Bo] R. E. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster, *Proc. Natl. Acad. Sci. USA* **83** (1986), 3068-3071.
- [Br] P. Bressler, Vertex algebroids II, arXiv: math.AG/0304115.
- [BrC] P. Bressler and A. Chervov, Courant algebroids, arXiv: hep-th/0212195.
- [Co] T. Courant, Dirac manifolds, *Trans. A.M.S.* **319** (1990), 631-661.
- [Di] L.A. Dickey, Soliton equations and Hamiltonian systems, *World Scientific* (1991).
- [DLiM] C. Dong, H.-S. Li and G. Mason, Vertex Lie algebra, vertex poisson algebras and vertex algebras, in: *Recent Developments in Infinite-Dimensional Lie Algebras and Conformal Field Theory*, Proceedings of an International Conference at University of Virginia, May 2000, Contemporary Math. **297** (2002), 69-96.

- [FB] D. Ben-Zvi and E. Frenkel, Vertex Algebras and Algebraic Curves, Mathematical Surveys and Monographs, **Vol. 88**, Amer. Math. Soc., Providence, 2001.
- [FHL] I. Frenkel, Y.-Z. Huang and J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, *Memoirs Amer. Math. Soc.* **Vol. 104**, no. 494, 1993.
- [GMS] V. Gorbounov, F. Malikov and V. Schechtman, Gerbes of chiral differential operators, II, Vertex Algebroids, *Invent. Math.* **155** (2004), no. 3, 605-680.
- [K] V. G. Kac, *Vertex Algebras for Beginners*, University Lecture Series **10**, Amer. Math. Soc., 1997.
- [LLi] J. Lepowsky and H.-S. Li, Introduction to Vertex Operator Algebras and Their Representations, *Progress in Math.*, **227**, Birkhäuser, Boston, 2003.
- [Li1] Representation theory and tensor product for vertex operator algebras, Ph.D. thesis, Rutgers University, 1994.
- [Li2] H.-S. Li, Vertex algebras and vertex Poisson algebras, *Commun. Contemp. Math.* **6** (2004), 61-110.
- [Li3] H.-S. Li, Abelianizing vertex algebras, *Commun. Math. Phys.* **259** (2005), 391-411.
- [LiY] H.-S. Li and G. Yamskulna, On certain vertex algebras and their modules associated with vertex algebroids, *J. Alg.* **283** (2005), 367-398.
- [LiuWX] Z.-J. Liu, A. Weinstein and P. Xu, Mainn triples for Lie bialgebroids, *J. Diff. Geometry* **45** (1997), 547-574.
- [LiuX] Z.-J. Liu and P. Xu, Dirac structures and dynamical r -matrices, math.DG/9903110.
- [P] M. Primc, Vertex algebras generated by Lie algebras, *J. Pure Appl. Alg.* **135** (1999), 253-293.
- [Ro] D. Roytenberg, Courant algebroids, derived brackets and even symplectic supermanifolds, Ph.D. thesis, UC Berkeley, 1999; math.DG/9910078.
- [RoW] D. Roytenberg and A. Weinstein, Courant algebroids and strongly homotopy Lie algebras, *Lett. Math. Phys.* **46** (1998), 81-93.